

Eur J Vasc Endovasc Surg 32, 286–293 (2006)

doi:10.1016/j.ejvs.2006.03.009, available online at <http://www.sciencedirect.com> on  SCIENCE @ DIRECT®

REVIEW

Can Statins Reduce Perioperative Morbidity and Mortality in Patients Undergoing Non-Cardiac Vascular Surgery?

K.I. Paraskevas,¹ C.D. Liapis,¹ G. Hamilton² and D.P. Mikhailidis^{3*}

¹Department of Vascular Surgery, Athens University Medical School, Athens, Greece,

²Academic Department of Surgery, and ³Department of Clinical Biochemistry, Royal Free and University College Medical School, London, UK

Aims. To determine whether statins can reduce perioperative morbidity and mortality in patients undergoing non-cardiac vascular surgery.

Methods. A search using Pubmed was performed to identify reports in English. The search terms were: “statins”, “perioperative morbidity”, “perioperative mortality” and “vascular surgery”. We excluded studies dealing with the effect of statins in cardiac surgery. Retrieved articles were manually searched.

Results. Current evidence shows that statins decrease perioperative morbidity and mortality in patients undergoing non-cardiac vascular surgery. Any benefit probably occurs soon (within a month) after initiating treatment.

Conclusions. Appropriately designed trials need to confirm the beneficial effect of perioperative statin therapy in various patient categories. The optimal duration and dose of perioperative statin therapy should be defined.

Keywords: Statins; Cardiovascular disease; Perioperative mortality; Vascular surgery.

Introduction

The prevalence of vascular complications is as high as 5.6% in non-cardiac operations.¹ Therefore, as many as 1 million surgical procedures may be complicated by perioperative vascular events every year.² Complications, such as atrial fibrillation,^{3–4} ventricular arrhythmias,⁵ pulmonary edema⁶ and myocardial infarction (MI),^{6–8} are the main causes of increased perioperative,^{7,8} as well as long-term,⁹ morbidity and mortality in patients undergoing major elective non-cardiac surgery. Postoperative myocardial ischaemic events are common in high-risk patients (e.g. those with coronary heart disease) undergoing non-cardiac surgery.¹⁰ Lee *et al.* analyzed the risk of major cardiac complications (MI, cardiogenic pulmonary edema, cardiac arrest and cardiac death) in 4315 patients

undergoing non-cardiac surgery.¹¹ The Lee index can predict major cardiac complications and cardiovascular death in patients undergoing non-cardiac surgery,¹² but its classification of procedures seems suboptimal.

MI is a cause of perioperative morbidity and mortality after vascular surgery^{2,13} that may remain unrecognised.^{11,14–16} Although the incidence of perioperative MI among patients undergoing major non-cardiac surgery is 2–3%, rates among those undergoing major vascular surgery can reach 34%.^{7,16} Endovascular surgery may be associated with reduced perioperative complications compared with open surgery, although this does not apply to late cardiac events.¹⁷ Perioperative MI is also associated with prolonged hospital stay and morbidity and mortality rates as high as 25–40%.^{18,19}

Various drugs may reduce vascular events in patients undergoing major elective vascular surgery. These include beta-blockers,^{7,8,20–25} calcium-channel blockers,²⁶ α_2 -adrenergic agonists,^{27–29} and aspirin.³⁰ However, no globally accepted guidelines have been formulated.³¹ It follows that patients undergoing elective vascular surgery are often underprotected.³²

*Corresponding author. Dr Dimitri P. Mikhailidis, MD, FFPM, FRCPATH, FRCP, Department of Clinical Biochemistry (Vascular Disease Prevention Clinics), Royal Free Hospital, Royal Free and University College Medical School, Pond Street, London NW3 2QG, UK.

E-mail address: mikhailidis@aol.com

Statins can stabilize coronary plaques,^{33–36} and exert anti-inflammatory antithrombotic, antiproliferative and leucocyte-adhesion inhibiting effects.^{37–44} Therefore, statins can prevent vascular complications,⁴⁵ possibly cost-effectively.⁴⁶

Methods

We searched PubMed for studies in English using the terms “statins”, “perioperative morbidity”, “perioperative mortality” and “vascular surgery”. We excluded articles dealing with cardiac surgery or percutaneous intervention. The relevant articles were manually searched for additional studies.

Results

Perioperative statin therapy and morbidity/mortality rates

Trials that investigated the effect of perioperative statin therapy on morbidity and mortality in patients undergoing non-cardiac vascular surgery are shown in Table 1.

In a retrospective case-control study including 2816 patients undergoing major non-cardiac vascular surgery (acute or elective abdominal aortic aneurysm repair, carotid endarterectomy [CEA], lower extremity revascularization),⁴⁷ perioperative statin use was associated with a > 4-fold reduced risk in perioperative mortality [adjusted odds ratio (OR) = 0.22, 95% confidence interval (CI) = 0.10–0.47]. This finding was supported by a prospective, randomized, placebo-controlled, double-blind trial assessing the effect of atorvastatin (20 mg/day) for 45 days on perioperative cardiovascular events after vascular surgery compared with placebo.⁴⁸ Vascular surgery was performed not earlier than 2 weeks after randomization (average 31 days after starting medication). All patients were followed for 6 months after surgery. Despite the small number of patients (50/group), a more than three-fold risk reduction was observed for statin users vs. placebo (8.0% vs. 26.0%, $p = 0.031$). The rate of combined end points (death from cardiac causes, nonfatal acute MI, ischemic stroke and unstable angina) was 9.1% in the atorvastatin group and 28.3% in the placebo group ($p = 0.030$).

The beneficial effect of statins and non-statin lipid-lowering agents was reported in a large, retrospective cohort study including 780 591 patients undergoing major non-cardiac surgery.⁴⁹ Of these, 65 399 patients underwent vascular surgery; 13 862 patients were

statin users. Statin use was associated with a significant ($p < 0.001$) reduction in perioperative mortality in patients undergoing not only vascular, but also any non-cardiac surgery.

A study [Statins for Risk Reduction in Surgery (StaRRS)] assessing the effect of statins on cardiac complications (death, MI, myocardial ischemia, ventricular tachyarrhythmias and acute congestive heart failure) in patients undergoing non-cardiac vascular surgery (CEA, aortic surgery, or lower extremity revascularization) supported the use of perioperative statin therapy.⁵⁰ Complications occurred in 157 of 1163 eligible hospitalizations. A multivariate model (accounting for age, gender, body mass index, type of operation, whether surgery was urgent or elective, diabetes mellitus and left ventricular dysfunction) showed that statins were beneficial (OR = 0.52, 95% CI = 0.35–0.76, $p = 0.001$).

Another trial included 1566 patients undergoing CEA (1440 patients) or CEA combined with coronary artery bypass grafting (CABG, 126 patients). Perioperative statin (20 ± 10 mg/day atorvastatin [332 patients], 20 ± 10 mg/day simvastatin [189 patients], 30 ± 20 mg/day pravastatin [91 patients], 30 ± 10 mg/day lovastatin [32 patients] or 30 ± 10 mg/day fluvastatin [13 patients]) use in 657 (42%) patients significantly reduced the incidence of cerebrovascular events and mortality.⁵¹ In multivariate analysis, adjusting for comorbidities associated with stroke (symptomatic carotid artery disease, chronic atrial fibrillation, hyperlipidemia, intraluminal shunt and patch graft use and combined CEA/CABG), statin use was associated with reduced odds of perioperative stroke (OR = 0.29, 95% CI = 0.14–0.61, $p < 0.05$).⁵¹ Similarly, in multivariate analysis of perioperative mortality, adjusting for comorbidities (percentage carotid stenosis, hypertension, chronic atrial fibrillation, coronary artery disease, congestive heart failure, chronic renal insufficiency, use of β -blockers, and combined CEA/CABG), statins reduced perioperative death (OR = 0.14, 95% CI = 0.03–0.62, $p < 0.05$).⁵¹ An observational study verified the protective effect of statins in patients undergoing CEA (Table 1).⁵² However, as the authors underlined, these results need to be confirmed in randomized controlled trials.

The long-term benefit of statins was reported in patients undergoing successful abdominal aortic aneurysm (AAA) surgery.^{53–55} A 6.3% risk for cardiac death and MI at 2 years,⁵⁴ and a 28% incidence for death due to cardiovascular causes at 8 years⁵⁵ following aortic operations have been reported. After a median follow-up of 4.7 years, patients using statins (154 of 510 followed-up patients) had a 2.5-fold reduction

Table 1. Trials assessing the effect of statin therapy on perioperative morbidity and mortality in patients undergoing non-cardiac vascular surgery

Study (Year)	Number of patients	Study Design	Statin used	Type of operation	Outcome
Poldermans <i>et al.</i> ⁴⁷ (2004)	2816	Retrospective case-controlled study including 2816 patients, 160 case subjects were identified who died because of any cause during surgery or during post-operative hospital stay <30 days. For each case subject, 2 controls were selected (one before and one after the case), i.e. 320 patients formed the control group.	NM	<ul style="list-style-type: none"> – Acute AAA repair: 38% – Elective AAA repair: 48% – CEA: 4% – Lower extremity revascularization: 11% 	Perioperative mortality was reduced 4.5 times compared to non-users (adjusted OR = 0.22, 95% CI = 0.10–0.47)
Durazzo <i>et al.</i> ⁴⁸ (2004)	100 50 statin-users and 50 placebo	Prospective, randomised, placebo-controlled, double blind trial.	Atorvastatin 20 mg/day or placebo for 45 days.	<ul style="list-style-type: none"> – AAA repair: 28% vs. 26%* – Infrainguinal arterial bypass: 18% vs. 22%* – CEA: 12% vs. 10%* – Amputation: 2% vs. 4%* – No operation: 12% vs. 8%* <p>* Percentages are given for patients on atorvastatin vs. placebo</p>	Overall rate of combined endpoints of death from cardiac causes, nonfatal acute MI, ischemic stroke and unstable angina was 8.0% vs. 26.0%, in the atorvastatin and placebo groups, respectively ($p = 0.031$).
Lindenauer <i>et al.</i> ⁴⁹ (2004)	For vascular surgery procedures: 65399 13862 statin users vs. 51537 non-users	Retrospective cohort study including 780591 patients undergoing major non-cardiac surgery (vascular, orthopedic, abdominal, thoracic, gynecologic, urologic, neurosurgical, otolaryngologic, transplant, trauma, and plastic and reconstructive)	NM	NM	For the whole study statin use was associated with a significant ($p < 0.001$) reduction in perioperative mortality (1595 users vs. 4158 non-users, $p < 0.001$). Results were not provided for each type of operation.
O'Neil-Callahan <i>et al.</i> ⁵⁰ (2005)	1163	Retrospective study of CEAs, aortic surgery (aorto-iliac bypass, AAA or dissection repair), or peripheral lower extremity revascularization procedures. A total of 1163 hospitalizations on 997 patients were included in the study.	NM	<ul style="list-style-type: none"> – CEA: 31.3% – Aortic Surgery (aorto-iliac bypass, AAA, dissection repair): 15.2% – Lower extremity revascularization: 53.5% 	Statin use associated with a significant reduction in the rate of complications in both univariate (OR = 0.56, 95% CI = 0.39–0.79, $p = 0.001$) and multi-variate (OR = 0.52, 95% CI = 0.35–0.76, $p = 0.001$) analyses.
McGirt <i>et al.</i> ⁵¹ (2005)	1566 657 statin users vs. 909 non-users (42% vs. 58%)	Retrospective study	Atorvastatin: 332 pts Simvastatin: 189 pts Pravastatin: 91 pts Lovastatin: 32 pts Fluvastatin: 13 pts	CEA	Statin use associated with a reduction in perioperative strokes (1.2% vs. 4.5%, $p < 0.01$), peri-operative TIAs (1.5% vs. 3.6%, $p < 0.01$), mortality (0.3% vs. 2.1%, $p < 0.01$), and the median (inter-quartile range) length of hospitalisation (2 days [2–5 days] vs. 3 days [2–7 days], $p < 0.05$)

Kennedy <i>et al.</i> ⁵² (2005)	3283 CEAs 1252 Asymptomatic (665 statin users), 2031 Symptomatic (815 statin users)	Observational Study	NM	CEA	When patients were matched to reduce the bias in estimation of potential treatment effect, statin use was associated with absolute reductions in in-hospital mortality rate of 0.9% ($p = 0.005$), 0.8% ($p = 0.0034$), and 0.9% ($p = 0.016$) using the kernal, stratified and radius matching methods, respectively. The reductions associated with statin use for in-hospital ischemic stroke or death rates were 1.7% ($p = 0.022$), 1.5% ($p = 0.055$) and 1.7% ($p = 0.038$), respectively
Kertai <i>et al.</i> ⁵³ (2004)	510 154 statin users (30%)	Retrospective study. Although 570 patients initially underwent AAA repair, 519 (91%) survived surgery for at least 30 days and a total of 510 patients were followed-up until the end of the study period [Median follow-up period: 4.7 years (interquartile range, 2.7 to 7.3 years)]	Simvastatin: 110 pts, 71% Atorvastatin: 27 pts, 18% Pravastatin: 16 pts, 10% Fluvastatin: 1 pt, 1%	AAA repair	After a median follow-up of 4.7 years (interquartile range, 2.7 to 7.3 years), statin use was shown to be associated with a 2.5-fold reduction in the risk of all-cause mortality (HR = 0.4; 95% CI: 0.3–0.5) and more than a threefold reduction in the risk of cardiovascular mortality (HR = 0.3; 95% CI: 0.2 to 0.6)
Parker Ward <i>et al.</i> ⁵⁷ (2005)	446 72 statin users vs. 374 non-users	Retrospective analysis of infrainguinal vascular bypass surgeries focusing on the association between preoperative statin and beta-blocker use, perioperative complications and length of hospital stay.	NM	Infrainguinal arterial bypass surgery	Statin use associated with fewer combined cardiovascular complications (6.9% vs. 20.1%, $p < 0.0008$) and a shorter length of stay (6.4 vs. 9.7 days, $p < 0.007$).
Abbruzzese <i>et al.</i> ⁵⁸ (2004)	172 88 statin users vs. 84 non-users	A total of 172 patients and 189 grafts were included in the study. Of these, 88 patients (94 grafts) were in the statin group and 84 patients (95 grafts) belonged to the control group.	– Results were provided for limbs, not patients: Simvastatin: 60 limbs, 64% Atorvastatin: 28 limbs, 30% Cerivastatin: 1 limb, 1% Lovastatin: 3 limbs, 3% Pravastatin: 2 limbs, 2%	Infrainguinal arterial bypass surgery using autogenous saphenous vein graft	– Perioperative morbidity and mortality did not differ between the 2 groups. – Two-year primary-revised ($94\% \pm 2\%$ vs. $83\% \pm 5\%$, $p < 0.02$) and secondary ($97\% \pm 2\%$ vs. $87\% \pm 4\%$, $p < 0.02$) graft patency rates were higher for the statin group compared with the control group.
Kertai <i>et al.</i> ⁵⁹ (2004)	570 162 statin users	Retrospective study	Simvastatin 71.6% Atorvastatin 14.8% Pravastatin 8.6% Fluvastatin 1.2% Cerivastatin 3.7%	AAA repair	Statin users had a 3-fold reduced risk of a composite end-point of 30-day perioperative mortality and nonfatal MI compared with non-users [6 (3.7%) vs. 45 (11.0%) pts, respectively, OR = 0.31, 95% CI = 0.13–0.74, $p = 0.01$]

NM = Not Mentioned, AAA = Abdominal Aortic Aneurysm, CEA = Carotid Endarterectomy, OR = Odds Ratio, CI = Confidence Interval, pts = patients, HR = Hazard Ratio, MI = myocardial infarction.

in all-cause mortality (hazard ratio [HR] = 0.4; 95% CI = 0.3–0.5) and more than a threefold reduction in the risk of vascular mortality (HR = 0.3; 95% CI: 0.2–0.6) compared with non-users (356 of 510 followed-up patients).⁵³ In the same study, use of β -blockers was also associated with an approximately 1.5-fold reduction in all-cause and vascular mortality compared with non-use thus verifying the results of another study,⁵¹ although an opposite opinion was also expressed.⁵⁶ The authors concluded that statin use in patients undergoing successful AAA surgery is associated with a significant reduction in all-cause and vascular mortality. Similar results regarding long-term survival were obtained from preoperative statin therapy in patients undergoing infrainguinal vascular bypass surgery⁵⁷ in addition to improved graft patency.⁵⁸

A beneficial effect from using both β -blockers and statins on perioperative (30-day) mortality and non-fatal MI in patients undergoing AAA surgery was demonstrated.⁵⁹ In multivariate analysis, patients receiving both medications showed a 4-fold and 3-fold reduction in perioperative mortality and nonfatal MI following AAA surgery, respectively, compared with patients receiving either statins (7.7% vs. 33%, OR = 0.24, 95%CI = 0.10–0.70, $p = 0.01$) or β -blockers (7.7% vs. 20%, OR = 0.24, 95%CI = 0.11–0.54, $p = 0.01$) alone. These results⁵⁹ suggest a benefit of combining statins and β -blockers in high-risk patients as defined by a revised cardiac index. The opinion that combined treatment for vascular risk factors results in increased benefit has been reported.^{60–61}

A prospective, randomized trial (the Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echo – IV [DECREASE-IV] Study) is under way.⁶² It will determine the impact of perioperative administration of a β -blocker (bisoprolol), a statin (fluvastatin), or both, on 30-day vascular events (including non-fatal MI or any death with a cardiovascular cause: cardiac arrest, MI, pulmonary embolus, stroke, hemorrhage, or death due to unknown causes) following major non-cardiac surgery.

Statins and peripheral arterial disease (PAD)

In patients with PAD, statins improve lower extremity blood flow and lower extremity function (as defined by ankle brachial index),^{63–64} as well as subjective symptoms of claudication (e.g. walking time).^{65–68} In addition, a retrospective review⁵⁷ of infrainguinal vascular bypass surgery in 446 patients showed that preoperative statin use was associated with fewer cardiovascular complications (6.9% vs. 20.1%; $p < 0.008$)

and a shorter in-hospital stay compared with non-users (6.4 vs. 9.7; $p = 0.007$). Statin use was a predictor of fewer preoperative combined vascular complications in multivariate analysis (OR = 0.36, 95% CI = 0.14–0.93, $p = 0.035$).

The results that derive from statin use⁵⁷ are better compared with the results that derive from perioperative β -blockade use in patients undergoing infrarenal vascular surgery.²⁴ In this double-blind, randomized, placebo-controlled trial the effect of perioperative β -blockade with metoprolol vs. placebo on patients undergoing infrarenal vascular surgery was investigated.²⁴ Metoprolol use did not reduce 30-day cardiovascular events (MIs, unstable angina, ventricular tachycardia, stroke) compared with placebo [Adjusted (for sex, age, baseline use of statins and planned aortic cross-clamping) RR = 0.87 (0.48–1.55)]. However, metoprolol significantly reduced the median (95% CI) time between surgery and discharge compared with placebo (9 [8–12] vs. 12 [9–19]) days (adjusted [for sex, age, baseline use of statins and planned use of aortic cross-clamping] HR = 1.71, 95% CI = 1.09–2.66, $p = 0.02$).

It has been suggested that statins exert a renoprotective effect in PAD and other patients.^{69–73} There is also evidence that serum creatinine and calculated creatinine clearance are indicators of postoperative mortality.⁷⁴ Therefore, the effect of statins on renal function following vascular surgery deserves investigation.

National guidelines regarding the use of statins in PAD were proposed.⁷⁵ Similar international guidelines could result in a widely accepted strategy.

Safety of Statin Use

Perioperative statin administration in high-risk patients undergoing major elective vascular surgery seems to be safe⁷⁶ and not associated with statin-related myopathy.^{77–81} A total of 885 patients (211 statin users) undergoing major elective vascular surgery (527 AAA repair operations and 358 lower extremity revascularization procedures) were studied. Although maximum serum creatine kinase activities were significantly higher in statin users compared with non-users (301 for statin users, range 16–13 377 U/L vs. 192 for non-users, range 8–30 390, $p = 0.003$) the difference became non-significant ($p = 0.142$) when corrected for length of surgery, cardiac risk factors and risk for myopathy. No muscular complaints or case of rhabdomyolysis was observed in either group. In addition, significantly fewer nonfatal MIs (15 vs. 81 and 6.7% vs. 10.8%; $p < 0.01$) and perioperative deaths

(5 vs. 35 and 2.1% vs. 3.9%; $p < 0.01$) occurred in statin users compared with non-users. The combined endpoint of perioperative death and MI occurred in significantly fewer statin users compared with non-users (8.8% vs. 14.4%; $p < 0.01$).

The Future

More evidence is required to confirm the benefit observed in the studies described above. However, the obvious ethical restrictions may mean that we will never see placebo-controlled trials. One option is to evaluate different targets and various lipid-lowering drugs in the perioperative period to establish the optimal choice. For example, the cholesterol lowering effect of statins can be extended by adding ezetimibe, a selective cholesterol transport inhibitor.⁸²

The low density lipoprotein cholesterol (LDL-C) target for patients with PAD needs to be established. For this high risk population it would be desirable to achieve the LDL-C target defined by the European and USA guidelines for this category (i.e. 2.5 and 2.6 mmol/l, respectively with an optional target of 1.8 mmol/l for very high risk patients).^{83–84}

Conclusions

There is growing evidence suggesting that statin use significantly reduces perioperative morbidity and mortality in patients undergoing non-cardiac vascular surgery, in a similar manner to that reported in cardiac surgery,^{85–86} as well as in renal,^{87–88} heart,^{89–90} and lung⁹¹ transplantation operations. However, optimal perioperative statin therapy has not been defined. Appropriately designed trials are needed to define optimal perioperative statin therapy in various categories of patients.

References

- 1 BADNER NH, KNILL RL, BROWN JE, NOVICK TV, GELB AW. Myocardial infarction after noncardiac surgery. *Anesthesiology* 1998;**88**: 572–578.
- 2 MANGANO DT. Perioperative cardiac morbidity. *Anesthesiology* 1990;**72**:153–184.
- 3 PASSMAN RS, GINGOLD DS, AMAR D *et al.* Prediction rule for atrial fibrillation after major noncardiac thoracic surgery. *Ann Thorac Surg* 2005;**79**:1698–1703.
- 4 CHRISTIANS KK, WU B, QUEBBEMAN EJ, BRASEL KJ. Postoperative atrial fibrillation in noncardiothoracic surgical patients. *Am J Surg* 2001;**182**:713–715.
- 5 AMAR D, ZHANG H, ROISTACHER N. The incidence and outcome of ventricular arrhythmias after noncardiac thoracic surgery. *Anesth Analg* 2002;**95**:537–543.
- 6 FLEISCHMANN KE, GOLDMAN L, YOUNG B, LEE TH. Association between cardiac and noncardiac complications in patients undergoing noncardiac surgery: outcomes and effects on length of stay. *Am J Med* 2003;**115**:515–520.
- 7 POLDERMANS D, BOERSMA E, BAX JJ *et al.* The effect of bisoprolol on perioperative mortality and myocardial infarction in high-risk patients undergoing vascular surgery. Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography Study Group. *N Engl J Med* 1999;**341**:1789–1794.
- 8 MANGANO DT, LAYUG EL, WALLACE A, TATEO I. Effect of atenolol on mortality and cardiovascular morbidity after noncardiac surgery. Multicenter Study of Perioperative Ischemia Research Group. *N Engl J Med* 1996;**335**:1713–1720.
- 9 FARKOUH ME, RIHAL CS, GERSH BJ *et al.* Influence of coronary heart disease on morbidity and mortality after lower extremity revascularization surgery: a population-based study in Olmsted County, Minnesota (1970–1987). *J Am Coll Cardiol* 1994;**24**: 1290–1296.
- 10 MANGANO DT, BROWNER WS, HOLLENBERG M, LONDON MJ, TATEO IM. Association of perioperative myocardial ischemia with cardiac morbidity and mortality in men undergoing noncardiac surgery. The Study of Perioperative Ischemia Research Group. *N Engl J Med* 1990;**323**:1781–1788.
- 11 LEE TH, MARCANTONIO ER, MANGIONE C *et al.* Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999;**100**:1043–1049.
- 12 BOERSMA E, KERTAI MD, SCHOUTEN O *et al.* Perioperative cardiovascular mortality in noncardiac surgery: validation of the Lee cardiac index. *Am J Med* 2005;**118**:1134–1141.
- 13 FLEISHER LA, EAGLE KA. Clinical practice. Lowering cardiac risk in noncardiac surgery. *N Engl J Med* 2001;**345**:1677–1682.
- 14 GRIMM JR RH, TILLINGHAST S, DANIELS K *et al.* Unrecognized myocardial infarction: experience in the Multiple Risk Factor Intervention Trial (MRFIT). *Circulation* 1987;**75**:II6–II8.
- 15 SIGURDSSON E, THORGEIRSSON G, SIGVALDASON H, SIGFUSSON N. Unrecognized myocardial infarction: epidemiology, clinical characteristics, and the prognostic role of angina pectoris. The Reykjavik Study. *Ann Intern Med* 1995;**122**:96–102.
- 16 SHEIFER SE, GERSH BJ, YANEZ III ND, ADES PA, BURKE GL, MANOLIO TA. Prevalence, predisposing factors, and prognosis of clinically unrecognized myocardial infarction in the elderly. *J Am Coll Cardiol* 2000;**35**:119–126.
- 17 SCHOUTEN O, VAN WANING VH, KERTAI MD *et al.* Perioperative and long-term cardiovascular outcomes in patients undergoing endovascular treatment compared with open vascular surgery for abdominal aortic aneurysm or iliofemoro-popliteal bypass. *Am J Cardiol* 2005;**96**:861–866.
- 18 LINDENAUER PK, FITZGERALD J, HOOPLE N, BENJAMIN EM. The potential preventability of postoperative myocardial infarction: underuse of perioperative β -adrenergic blockade. *Arch Intern Med* 2004;**164**:762–766.
- 19 NETTLEMAN MD, BANITT L, BARRY W, AWAN I, GORDON EEI. Predictors of survival and the role of gender in postoperative myocardial infarction. *Am J Med* 1997;**103**:357–362.
- 20 POLDERMANS D, BOERSMA E, BAX JJ *et al.* Bisoprolol reduces cardiac death and myocardial infarction in high-risk patients as long as 2 years after successful major vascular surgery. *Eur Heart J* 2001;**22**: 1353–1358.
- 21 EAGLE KA, BERGER PB, CALKINS H *et al.* ACC/AHA guideline update for perioperative cardiovascular evaluation for noncardiac surgery – executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1996 Guidelines on Perioperative Cardiac Evaluation for Noncardiac Surgery). *J Am Coll Cardiol* 2002;**39**:542–553.
- 22 GRAYBURN PA, HILLIS LD. Cardiac events in patients undergoing noncardiac surgery: shifting the paradigm from noninvasive risk stratification to therapy. *Ann Intern Med* 2003;**138**:506–511.
- 23 WALLACE A, LAYUG B, TATEO I *et al.* Prophylactic atenolol reduces postoperative myocardial ischemia. *Anesthesiology* 1998;**88**:7–17.
- 24 BRADY AR, GIBBS JS, GREENHALGH RM, POWELL JT, SYDES MR; POBBLE trial investigators. Perioperative beta-blockade (POBBLE) for patients undergoing infrarenal vascular

- surgery: results of a randomized double-blind controlled trial. *J Vasc Surg* 2005;**41**:602–609.
- 25 KERTAI MD, BOERSMA E, BAX JJ *et al.* Dutch Echocardiographic Cardiac Risk Evaluation Applying Stress Echocardiography (DECREASE) Study Group. Optimizing long-term cardiac management after major vascular surgery: Role of beta-blocker therapy, clinical characteristics, and dobutamine stress echocardiography to optimize long-term cardiac management after major vascular surgery. *Arch Intern Med* 2003;**163**:2230–2235.
 - 26 WIJEYSUNDERA DN, BEATTIE WS. Calcium channel blockers for reducing cardiac morbidity and mortality after noncardiac surgery: a meta-analysis. *Anest Analg* 2003;**97**:634–641.
 - 27 OLIVER MF, GOLDMAN L, JULIAN DG, HOLME I. Effect of mivazerol on perioperative cardiac complications during non-cardiac surgery in patients with coronary heart disease: the European Mivazerol Trial (EMIT). *Anesthesiology* 1999;**91**:951–961.
 - 28 WALLACE AW, GALINDEZ D, SALAHIEH A *et al.* Effect of clonidine on cardiovascular morbidity and mortality after noncardiac surgery. *Anesthesiology* 2004;**101**:284–293.
 - 29 NISHINA K, MIKAWA K, UESUGI T *et al.* Efficacy of clonidine for prevention of perioperative myocardial ischemia: a critical appraisal and meta-analysis of the literature. *Anesthesiology* 2002;**96**:323–329.
 - 30 ROBLESS P, MIKHAILIDIS DP, STANSBY G. Systematic review of antiplatelet therapy for the prevention of myocardial infarction, stroke or vascular death in patients with peripheral arterial disease. *Br J Surg* 2001;**88**:787–800.
 - 31 DEVEREAX PJ, GOLDMAN L, YUSUF S, GILBERT K, LESLIE K, GUYATT GH. Surveillance and prevention of major perioperative ischemic events in patients undergoing noncardiac injury: a review. *CMAJ* 2005;**173**:779–788.
 - 32 HENKE PK, BLACKBURN S, PROCTOR MC *et al.* Patients undergoing infrainguinal bypass to treat atherosclerotic vascular disease are underprescribed cardioprotective medications: effect on graft patency, limb salvage, and mortality. *J Vasc Surg* 2004;**39**:357–365.
 - 33 DAWOOD MM, GUTPA DK, SOUTHERN J, WALLA A, ATKINSON JB, EAGLE KA. Pathology of fatal perioperative myocardial infarction: implications regarding pathophysiology and prevention. *Int J Cardiol* 1996;**57**:37–44.
 - 34 MORISHITA R, TOMITA N, OGIHARA T. HMG-Co A reductase inhibitors in the treatment of cardiovascular diseases: stabilization of coronary artery plaque. *Curr Drug Targets* 2002;**3**:379–385.
 - 35 IKEDA U, SHIMADA K. Pleiotropic effects of statins on the vascular tissue. *Curr Drug Targets Cardiovasc Haematol Disord* 2001;**1**:51–58.
 - 36 LUO JD, CHEN AF. Perspectives on the cardioprotective effects of statins. *Curr Med Chem* 2003;**10**:1593–1601.
 - 37 TAKEMOTO M, LIAO JK. Pleiotropic effects of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Arterioscler Thromb Vasc Biol* 2001;**21**:1712–1719.
 - 38 VAN HAELEST PL, VAN DOORMAAL JJ, MAY JF, GANS RO, CRIJNS HJ, COVEN TERVAERT JW. Secondary prevention with fluvastatin decreases levels of adhesion molecules, neopterin and C-reactive protein. *Eur J Intern Med* 2001;**12**:503–509.
 - 39 LAUFS U, LIAO JK. Direct vascular effects of HMG-CoA reductase inhibitors. *Trends Cardiovasc Med* 2000;**10**:143–148.
 - 40 LIBBY P, RIDKER PM, MASERI A. Inflammation and atherosclerosis. *Circulation* 2002;**105**:1135–1143.
 - 41 LAUFS U, LA FATA V, PLUTZKY J *et al.* Upregulation of endothelial nitric oxide synthase by HMG CoA reductase inhibitors. *Circulation* 1998;**97**:1129–1135.
 - 42 ALBERT MA, DANIELSON E, RIFAI N, RIDKER PM; PRINCE Investigators. Effect of statin therapy on C-reactive protein levels: the pravastatin inflammation/CRP evaluation (PRINCE): a randomized trial and cohort study. *JAMA* 2001;**286**:64–70.
 - 43 RIDKER PM, RIFAI N, PFEFFER MA *et al.* Inflammation, pravastatin, and the risk of coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events (CARE) Investigators. *Circulation* 1998;**98**:839–844.
 - 44 MILIONIS HJ, ELISAF MS, MIKHAILIDIS DP. The effect of lipid-regulating therapy on haemostatic parameters. *Curr Pharm Design* 2003;**9**:2425–2443.
 - 45 BICCARD BM, SEAR JW, FOEX P. Statin therapy: a potentially useful peri-operative intervention in patients with cardiovascular disease. *Anaesthesia* 2005;**60**:1106–1114.
 - 46 BICCARD BM, SEAR JW, FOEX P. The pharmacoeconomics of peri-operative statin therapy. *Anaesthesia* 2005;**60**:1059–1063.
 - 47 POLDERMANS D, BAX JJ, KERTAI MD *et al.* Statins are associated with a reduced incidence of perioperative mortality in patients undergoing major noncardiac vascular surgery. *Circulation* 2003;**107**:1848–1851.
 - 48 DURAZZO AE, MACHADO FS, IKEOKA DT *et al.* Reduction in cardiovascular events after vascular surgery with atorvastatin: a randomized trial. *J Vasc Surg* 2004;**39**:967–976.
 - 49 LINDENAUER PK, PEKOW P, WANG K, GUTIERREZ B, BENJAMIN EM. Lipid-lowering therapy and in-hospital mortality following major noncardiac surgery. *JAMA* 2004;**291**:2092–2099.
 - 50 O'NEIL-CALLAHAN K, KATSIMAGLIS G, TEPPER MR *et al.* Statins decrease perioperative cardiac complications in patients undergoing noncardiac vascular surgery. *J Am Coll Cardiol* 2005;**45**:336–342.
 - 51 MCGIRT MJ, PERLER BA, BROOKE BS *et al.* 3-Hydroxy-3-methylglutaryl coenzyme A reductase inhibitors reduce the risk of perioperative stroke and mortality after carotid endarterectomy. *J Vasc Surg* 2005;**42**:829–836.
 - 52 KENNEDY J, QUAN H, BUCHAN AM, GHALI WA, FEASBY TE. Statins are associated with better outcomes after carotid endarterectomy in symptomatic patients. *Stroke* 2005;**36**:2072–2076.
 - 53 KERTAI MD, BOERSMA E, WESTERHOUT CM *et al.* Association between long-term statin use and mortality after successful abdominal aortic aneurysm surgery. *Am J Med* 2004;**116**:96–103.
 - 54 KRUPSKI WC, LAYUG EL, REILLY LM, RAPP JH, MANGANO DT. Comparison of cardiac morbidity rates between aortic and infrainguinal operations: two-year follow-up. Study of Perioperative Ischemia Research Group. *J Vasc Surg* 1993;**18**:609–615.
 - 55 United Kingdom Small Aneurysm Trial Participants. Long-term outcomes of immediate repair compared with surveillance of small abdominal aortic aneurysms. *N Engl J Med* 2002;**346**:1445–1452.
 - 56 BOERSMA E, POLDERMANS D, BAX JJ *et al.* Predictors of cardiac events after major vascular surgery: role of clinical characteristics, dobutamine echocardiography, and beta-blocker therapy. *JAMA* 2001;**285**:1865–1873.
 - 57 PARKER WARD R, LEEPER NJ, KIRKPATRICK JN, LANG RM, SORRENTINO MJ, WILLIAMS KA. The effect of preoperative statin therapy on cardiovascular outcomes in patients undergoing infrainguinal vascular surgery. *Int J Cardiol* 2005;**104**:264–268.
 - 58 ABBRUZZESE TA, HAVENS J, BELKIN M *et al.* Statin therapy is associated with improved patency of autogenous infrainguinal bypass grafts. *J Vasc Surg* 2004;**39**:1178–1185.
 - 59 KERTAI MD, BOERSMA E, WESTERHOUT CM *et al.* A combination of statins and beta-blockers is independently associated with a reduction in the incidence of perioperative mortality and nonfatal myocardial infarction in patients undergoing abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 2004;**28**:343–352.
 - 60 ATHYROS VG, MIKHAILIDIS DP, PAPAGEORGIOU AA *et al.*; GREACE Study Collaborative Group. Effect of statins and ACE inhibitors alone and in combination on clinical outcome in patients with coronary heart disease. *J Hum Hypertens* 2004;**18**:781–788.
 - 61 ATHYROS VG, MIKHAILIDIS DP, PAPAGEORGIOU AA *et al.* Effect of statins and aspirin alone and in combination on clinical outcome in dyslipidemic patients with coronary heart disease. A subgroup analysis of the GREACE study. *Platelets* 2005;**16**:65–71.
 - 62 SCHOUTEN O, POLDERMANS D, VISSER L *et al.* Fluvastatin and bisoprolol for the reduction of perioperative cardiac mortality and morbidity in high-risk patients undergoing non-cardiac surgery: rationale and design of the DECREASE-IV study. *Am Heart J* 2004;**148**:1047–1052.
 - 63 MCDERMOTT MM, CRIQUI MH, LIU K *et al.* Lower ankle/brachial index, as calculated by averaging the dorsalis pedis and posterior tibial artery pressures, and association with leg functioning in peripheral arterial disease. *J Vasc Surg* 2000;**32**:1164–1171.

- 64 McDERMOTT MM, GREENLAND P, LIU K *et al*. The ankle brachial index is associated with leg function and physical activity: the Walking and Leg Circulation Study. *Ann Intern Med* 2002;**136**: 873–883.
- 65 MONDILLO S, BALLO P, BARBATI R *et al*. Effects of simvastatin on walking performance and symptoms of intermittent claudication in hypercholesterolemic patients with peripheral vascular disease. *Am J Med* 2003;**114**:359–364.
- 66 McDERMOTT MM, GURALNIK JM, GREENLAND P *et al*. Statin use and leg functioning in patients with and without lower-extremity peripheral arterial disease. *Circulation* 2003;**107**:757–761.
- 67 MOHLER III ER, HIATT WR, CREAGER MA. Cholesterol reduction with atorvastatin improves walking distance in patients with peripheral arterial disease. *Circulation* 2003;**108**:1481–1486.
- 68 DASKALOPOULOU SS, DASKALOPOULOS ME, LIAPIS CD, MIKHAILIDIS DP. Peripheral arterial disease: a missed opportunity to administer statins so as to reduce cardiac morbidity and mortality. *Curr Med Chem* 2005;**12**:443–452.
- 69 YOUSSEF F, SEIFALIAN AM, JAGROOP A *et al*. The early effect of lipid-lowering treatment on carotid and femoral intima media thickness (IMT). *Eur J Vasc Endovasc Surg* 2002;**23**:358–364.
- 70 ATHYROS VG, ELISAF M, PAPAGEORGIOU AA *et al*; GREACE Study Collaborative Group. Effect of statins versus untreated dyslipidemia on serum uric acid levels in patients with coronary heart disease: a subgroup analysis of the GREACE Atorvastatin and Coronary-heart-disease Evaluation (GREACE) study. *Am J Kidney Dis* 2004;**43**:589–599.
- 71 ATHYROS VG, MIKHAILIDIS DP, PAPAGEORGIOU AA *et al*. The effects of statins versus untreated dyslipidemia on renal function in patients with coronary heart disease. A subgroup analysis of the GREACE Atorvastatin and Coronary-heart-disease Evaluation (GREACE) study. *J Clin Pathol* 2004;**57**:728–734.
- 72 ELISAF M, MIKHAILIDIS DP. Statins and renal function. *Angiology* 2002;**53**:493–502.
- 73 ALNAEB ME, YOUSSEF F, MIKHAILIDIS DP, HAMILTON G. Short-term lipid-lowering treatment with atorvastatin improves renal function but not renal blood flow indices in patients with peripheral arterial disease. *Angiology* 2006;**57**:65–71.
- 74 KERTAI MD, BOERSMA E, BAX JJ, VAN DER MEIRACKER AH, VAN URK H, ROELANDT JR *et al*. Comparison between serum creatinine clearance for the prediction of postoperative mortality in patients undergoing major vascular surgery. *Clin Nephrol* 2003;**59**:17–23.
- 75 VAHL AC, REEKERS JA. The guideline 'Diagnosis and treatment of peripheral artery disease of the lower extremities' of The Netherlands Surgical Society. *Ned Tijdschr Geneesk* 2005;**149**:1670–1674.
- 76 SCHOUTEN O, KERTAI MD, BAX JJ *et al*. Safety of perioperative statin use in high-risk patients undergoing major vascular surgery. *Am J Cardiol* 2005;**95**:658–660.
- 77 THOMPSON PD, CLARKSON P, KARAS RH. Statin-associated myopathy. *JAMA* 2003;**289**:1681–1690.
- 78 BELLOSTA S, PAOLETTI R, CORSINI A. Safety of statins: focus on clinical pharmacokinetics and drug interactions. *Circulation* 2004;**109**(23 Suppl. 1):III50–III57.
- 79 ROSENSON RS. Current overview of statin-induced myopathy. *Am J Med* 2004;**116**:408–416.
- 80 PASTERNAK RC, SMITH Jr SC, BAIREY-MERZ CN *et al*. ACC/AHA/NHLBI clinical advisory on the use and safety of statins. *Circulation* 2002;**106**:1024–1028.
- 81 MOOSMANN B, BEHL C. Selenoprotein synthesis and side-effects of statins. *Lancet* 2004;**363**:892–894.
- 82 MIKHAILIDIS DP, WIERZBICKI AS, DASKALOPOULOU SS *et al*. The use of ezetimibe in achieving low density lipoprotein lowering goals in clinical practice: position statement of a United Kingdom consensus panel. *Curr Med Res Opin* 2005;**21**:959–969.
- 83 GRUNDY SM, CLEEMAN JJ, MERZ CN *et al*. National heart, lung, and blood institute; American college of cardiology foundation; American heart foundation. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation* 2004;**110**:227–239.
- 84 DE BACKER G, AMBROSIONI E, BORCH-JOHNSEN K *et al*. European Society of Cardiology. American Heart Association. American College of Cardiology. European guidelines on cardiovascular disease prevention in clinical practice. Third Joint Task Force of European and other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of eight societies and by invited experts). *Atherosclerosis* 2004;**173**: 381–391.
- 85 PASCERI V, PATTI G, NUSCA A, PRISTIPINO C, RICHICHI G, DI SCIASCIO G; ARMYDA Investigators. Randomized trial of atorvastatin for reduction of myocardial damage during coronary intervention: results from the ARMYDA (Atorvastatin for Reduction of Myocardial Damage during Angioplasty) study. *Circulation* 2004;**110**:674–678.
- 86 LAZAR HL. Role of statin therapy in the coronary bypass surgery. *Ann Thorac Surg* 2004;**78**:730–740.
- 87 COSIO FG, PESAVENTO TE, PELLETIER RP *et al*. Patient survival after renal transplantation III: the effects of statins. *Am J Kidney Dis* 2002;**32**:713–716.
- 88 LENTINE KL, BRENNAN DC. Statin use after renal transplantation: a systematic quality review of trial-based evidence. *Nephrol Dial Transplant* 2004;**19**:2378–2386.
- 89 WU AH, BALLANTYNE CM, SHORT BC *et al*. Statin use and risks of death or fatal rejection in the Heart Transplant Lipid Registry. *Am J Cardiol* 2005;**95**:367–372.
- 90 WENKE K, MEISER B, THIERY J, REICHART B. Impact of simvastatin therapy after heart transplantation: an 11-year prospective evaluation. *Herz* 2005;**30**:431–432.
- 91 JOHNSON BA, IACONO AT, ZEEVI A, MCCURRY KR, DUNCAN SR. Statin use is associated with improved function and survival of lung allografts. *Am J Respir Crit Care Med* 2003;**167**:1271–1278.

Accepted 26 March 2006

Available online 11 May 2006